

## A) Scientific Abstract

Several melanoma antigens have been identified and include MAGE 1 and 3, gp100, tyrosinase and MART-1/Melan-A. MART-1 is widely expressed in melanoma and is recognized by the majority of HLA-A2 restricted cytotoxic T lymphocytes (CTL) generated from tumor infiltrating lymphocytes (TIL). An immunodominant peptide derived from the MART-1 118 amino acid protein (13kD) designated MART<sub>27-35</sub> is expressed by HLA-A2.1 melanoma cells<sup>1</sup>. In addition, peptide MART<sub>27-35</sub> can be presented by HLA-A2.2, 2.4, 2.5, 2.6, 2.9 and 69.1 major histocompatibility complex subtypes<sup>2</sup>.

Peripheral blood mononuclear cells (PBMC) can be differentiated into potent antigen-presenting cells (APC) with dendritic cell (DC) properties following the method of Romani *et al*<sup>3</sup>, and these cultured DC have been shown to be superior to other APC for the generation of specific immune effectors. Transfection of a tumor antigen gene into DC would enable a continuous processing and presentation of antigenic peptides in the context of the patient's HLA subtype<sup>4,5,6</sup>. Genetically engineered DC are able to induce specific CTLs with fewer *in vitro* stimulations than when DC are exogenously pulsed with peptides<sup>4,6</sup>. We have constructed a replicant-deficient, E1-deleted adenoviral vector carrying the MART-1 gene (AdVMART1). Transduction of *in vitro* cultured HLA-A2.1 and A2.4 DC with AdVMART1 allows these cells to process and present appropriate MART-1 peptides capable of stimulating MART-1-specific T cells<sup>7,8</sup>.

We have also developed a preclinical model using murine tumor cells expressing the MART-1 gene<sup>9</sup>. In this model, AdVMART1-transduced murine DC elicit a strong systemic immune response capable of treating established tumor implants. The observed response is MHC class I-restricted and MART-1-specific as assessed *in vitro* and *in vivo*. Our murine model suggests that the ability to generate CTL with MART-1-expressing human DC will have therapeutic benefit in the treatment of cancer subjects.

1. Kawakami Y. *et al*. Cloning of the gene coding for a shared melanoma antigen recognized by autologous T cells infiltrating into tumor. *Proc Natl Acad Sci USA* 91: 3515, 1994.
2. Fleischhauer K. *et al*. Multiple HLA-A alleles can present an immunodominant peptide of the human melanoma antigen Melan-A/MART-1 to a peptide-specific HLA-A\*0201+ cytotoxic T cell line. *J Immunol* 157: 787, 1996.
3. Yee C. *et al*. Isolation of tyrosinase-specific CD8+ and CD4+ T cell clones from the peripheral blood of melanoma patients following *in vitro* stimulation with recombinant vaccinia virus. *J Immunol* 157: 4079, 1996.
4. Romani N. *et al*. Proliferating dendritic cell progenitors in human blood. *J. Exp Med* 180: 83, 1994.
5. Reeves M.E. *et al*. Retroviral transduction of human dendritic cells with a tumor-associated antigen gene. *Cancer Res* 56: 5672, 1996.
6. Marincola F.M. *et al*. Differential anti-MART-1/MelanA CTL activity in peripheral blood of HLA-A2 melanoma patients in comparison to healthy donors: evidence of *in vivo* priming by tumor cells. *J Immunother* 19: 266, 1996.
7. Butterfield L.H. *et al*. Generation of melanoma-specific cytotoxic T lymphocytes by dendritic cells transduced with a MART-1 adenovirus. *J Immunol* 1998 (in press).
8. Perez-Diez A. *et al*. Generation of CD8+ and CD4+ T cell responses to dendritic cells genetically engineered to express the MART-1/Melan-a gene. *Cancer Res* 1998 (in press).
9. Ribas A. *et al*. Genetic immunization to the melanoma antigen MART-1/Melan-A using recombinant adenovirus-transduced murine dendritic cells. *Cancer Res* 57: 2865, 1997.